

Morse theory for group presentations

Ximena Fernández

Categories & Topology. Mathematical Congress of the Americas 2021

Swansea University

EPSRC Centre for Topological Data Analysis



The Andrews–Curtis conjecture

Group presentations \longleftrightarrow Groups

$$\mathcal{P} = \langle x_1, x_2, \dots, x_n \mid r_1, r_2, \dots, r_m \rangle \quad G = F(x_1, x_2, \dots, x_n) / N(r_1, r_2, \dots, r_m)$$

Group presentations \longleftrightarrow Groups

$$\mathcal{P} = \langle x_1, x_2, \dots, x_n \mid r_1, r_2, \dots, r_m \rangle \quad G = F(x_1, x_2, \dots, x_n) / N(r_1, r_2, \dots, r_m)$$

Theorem [Novikov-Boone, 1955-1958]

There exists a finitely presented group G such that the **word problem** for G is **undecidable**.

Theorem [Adian-Rabin, 1957-1958]

The **isomorphism problem** in groups is **undecidable**.

Tietze's transformations

Theorem [Tietze, 1908]

Any finite presentation $\mathcal{P} = \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ of a group G can be transformed into any other presentation of the same group by a finite sequence of the following **operations**:

1. replace some relator r_i by r_i^{-1} ;
2. replace some relator r_i by $r_i r_j$ for some $j \neq i$;
3. replace some relator r_i by a conjugate $w r_i w^{-1}$ for some w in the free group $F(x_1, x_2, \dots, x_n)$;
4. replace each relator r_i by $\phi(r_i)$ where ϕ is an automorphism of $F(x_1, x_2, \dots, x_n)$;
5. add a generator x_{n+1} and a relator r_{m+1} that coincides with x_{n+1} , or the inverse of this operation;
6. add a relator 1, or the inverse of this operation.

Tietze's transformations

Theorem [Tietze, 1908]

Any finite presentation $\mathcal{P} = \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ of a group G can be transformed into any other presentation of the same group by a finite sequence of the following **operations**:

- Q^{**} {
1. replace some relator r_i by r_i^{-1} ;
 2. replace some relator r_i by $r_i r_j$ for some $j \neq i$;
 3. replace some relator r_i by a conjugate $w r_i w^{-1}$ for some w in the free group $F(x_1, x_2, \dots, x_n)$;
 4. replace each relator r_i by $\phi(r_i)$ where ϕ is an automorphism of $F(x_1, x_2, \dots, x_n)$;
 5. add a generator x_{n+1} and a relator r_{m+1} that coincides with x_{n+1} , or the inverse of this operation;
 6. add a relator 1, or the inverse of this operation.

The Andrews–Curtis conjecture

Conjecture [Andrews & Curtis, 1965]

Any balanced presentation $\mathcal{P} = \langle x_1, \dots, x_n \mid r_1, \dots, r_n \rangle$ of the trivial group can be transformed into $\langle \mid \rangle$ by a finite sequence of Q^{**} -transformations.

Conjecture [Andrews & Curtis, 1965]

Any balanced presentation $\mathcal{P} = \langle x_1, \dots, x_n \mid r_1, \dots, r_n \rangle$ of the trivial group can be transformed into $\langle \mid \rangle$ by a finite sequence of Q^{**} -transformations.

Potential counterexamples

- $\mathcal{P} = \langle x, y \mid xyx = yxy, x^n = y^{n+1} \rangle, n \geq 2$ [Akbulut & Kirby, 1985]
- $\mathcal{P} = \langle x, y \mid x^{-1}y^n x = y^{n+1}, x = y^{-1}xyx^{-1} \rangle, n \geq 2$ [Miller & Schupp, 1999]
- $\mathcal{P} = \langle x, y \mid x = [x^m, y^n], y = [y^p, x^q] \rangle, n, m, p, q \in \mathbb{Z}$ [Gordon, 1984]

Group presentations \longleftrightarrow CW-complexes of dim 2

Group presentations \longleftrightarrow CW-complexes of dim 2


$\mathcal{P} \longrightarrow K_{\mathcal{P}}$



Group presentations \longleftrightarrow CW-complexes of dim 2


$\mathcal{P}_K \longleftarrow K$

$\langle x_1 \dots, x_9 \mid x_7, x_6^{-1} x_7^{-1} x_3,$
 $x_3^{-1} x_9, x_1^{-1} x_3^{-1} x_{10}, x_4^{-1} x_5,$
 $x_5 x_6^{-1}, x_7, x_6 x_7^{-1} x_8, x_8^{-1} x_9,$
 $x_6 x_9^{-1} \rangle$








Topological point of view



Topological point of view






Conjecture [Andrews–Curtis, 1965]

Any contractible 2-complex 3-deforms to a point.

Discrete Morse theory

Discrete Morse theory

Let K be a **regular** CW-complex.




Discrete Morse theory

Let K be a **regular** CW-complex.

- A map $f : K \rightarrow \mathbb{R}$ is a **discrete Morse function** if for every cell e^n in K :

$$\#\{e^n > e^{n-1} : f(e^n) \leq f(e^{n-1})\} \leq 1 \quad \text{and} \quad \#\{e^n < e^{n+1} : f(e^n) \geq f(e^{n+1})\} \leq 1.$$




Discrete Morse theory

Let K be a **regular** CW-complex.

- A map $f : K \rightarrow \mathbb{R}$ is a **discrete Morse function** if for every cell e^n in K :

$$\#\{e^n > e^{n-1} : f(e^n) \leq f(e^{n-1})\} \leq 1 \quad \text{and} \quad \#\{e^n < e^{n+1} : f(e^n) \geq f(e^{n+1})\} \leq 1.$$




Discrete Morse theory

Let K be a **regular** CW-complex.

- A map $f : K \rightarrow \mathbb{R}$ is a **discrete Morse function** if for every cell e^n in K :

$$\#\{e^n > e^{n-1} : f(e^n) \leq f(e^{n-1})\} \leq 1 \quad \text{and} \quad \#\{e^n < e^{n+1} : f(e^n) \geq f(e^{n+1})\} \leq 1.$$

- An n -cell $e^n \in K$ is a **critical cell of index n** if the values of f in every face and coface of e^n increase with dimension.




Discrete Morse theory

Let K be a **regular** CW-complex.

- A map $f : K \rightarrow \mathbb{R}$ is a **discrete Morse function** if for every cell e^n in K :

$$\#\{e^n > e^{n-1} : f(e^n) \leq f(e^{n-1})\} \leq 1 \quad \text{and} \quad \#\{e^n < e^{n+1} : f(e^n) \geq f(e^{n+1})\} \leq 1.$$

- An n -cell $e^n \in K$ is a **critical cell of index n** if the values of f in every face and coface of e^n increase with dimension.




Discrete Morse theory

Let K be a **regular** CW-complex.

- A map $f : K \rightarrow \mathbb{R}$ is a **discrete Morse function** if for every cell e^n in K :

$$\#\{e^n > e^{n-1} : f(e^n) \leq f(e^{n-1})\} \leq 1 \quad \text{and} \quad \#\{e^n < e^{n+1} : f(e^n) \geq f(e^{n+1})\} \leq 1.$$

- An n -cell $e^n \in K$ is a **critical cell of index n** if the values of f in every face and coface of e^n increase with dimension.








Theorem [Forman, 1995]

Let K be a regular CW-complex and let $f : K \rightarrow \mathbb{R}$ be a discrete Morse function. Then K is homotopy equivalent to a CW-complex K_M with exactly one cell of dimension k for every critical cell of index k .

Morse theory for cell complexes

Theorem [Forman, 1995]

Let K be a regular CW-complex and let $f : K \rightarrow \mathbb{R}$ be a discrete Morse function. Then K is homotopy equivalent to a CW-complex $K_{\mathcal{M}}$ with exactly one cell of dimension k for every critical cell of index k .



?

Goals:


Given K a regular CW-complex k of dimension n and a discrete Morse function $f : K \rightarrow \mathbb{R}$, we aim to:

- (re)construct the Morse complex $K_{\mathcal{M}}$,
- recover information about the simple homotopy type of K and moreover its $(n + 1)$ -deformation class.

Morse theory and collapses

Level subcomplex



$$K(c) = \bigcup_{\substack{e \in K \\ f(e) \leq c}} \bar{e}$$




$K(14)$

$K(13)$

$K(12)$




$K(6)$



$K(5)$



$K(4)$



$K(3)$



$K(1)$

Lema [F.]

Let K be a regular CW-complex.

Then, $f : K \rightarrow M$ is a discrete Morse function with critical cells C if and only if there exist a **sequence of subcomplexes** of K

$$K_0 \subseteq L_1 \subseteq K_1 \cdots \subseteq K_{N-1} \subseteq L_{N-1} \subseteq K_N = K$$

such that $K_j \searrow L_j$ for all $1 \leq j \leq N$ and the set of cells of K that was not collapsed in any of the collapses $K_j \searrow L_j$ is equal to C .


Morse theory and Whitehead deformations

Lema

Let K be a CW-complex of dimension $\leq n$. Let $\varphi : \partial D^n \rightarrow K$ be the attaching map of an n -cell e^n . If $K \searrow L$, then

$$K \cup e^n \xrightarrow{n+1} L \cup \tilde{e}^n$$

where the attaching map $\tilde{\varphi} : \partial D^n \rightarrow L$ of \tilde{e}^n is defined as $\tilde{\varphi} = r\varphi$ with $r : K \rightarrow L$ the canonical strong deformation retract induced by the collapse $K \searrow L$.



Morse theory and Whitehead deformations

Lema

Let K be a CW-complex of dimension $\leq n$. Let $\varphi : \partial D^n \rightarrow K$ be the attaching map of an n -cell e^n . If $K \searrow L$, then

$$K \cup e^n \xrightarrow{n+1} L \cup \tilde{e}^n$$

where the attaching map $\tilde{\varphi} : \partial D^n \rightarrow L$ of \tilde{e}^n is defined as $\tilde{\varphi} = r\varphi$ with $r : K \rightarrow L$ the canonical strong deformation retract induced by the collapse $K \searrow L$.

Definition

We say that there is an **internal collapse** from $K \cup e^n$ to $L \cup \tilde{e}^n$.

Theorem [F.]

Let K be a CW-complex on dimension n . Let

$$\emptyset = K_{-1} \subseteq L_0 \subseteq K_0 \subseteq L_1 \subseteq K_1 \cdots \subseteq L_N \subseteq K_N = K$$

be a sequence of subcomplexes of L such that $K_j \searrow L_j$ for all $j = 0, 1, \dots, N$. If

$$L_j = K_{j-1} \cup \bigcup_{i=1}^{d_j} e_i^j, \text{ then}$$

$$K \begin{array}{c} \nearrow \\ \searrow \end{array} \begin{array}{c} n+1 \\ \searrow \end{array} L_0 \cup \bigcup_{j=0}^N \bigcup_{i=1}^{d_j} \tilde{e}_i^j.*$$

*Here, the attaching maps of the cells \tilde{e}_i^j can be explicitly reconstructed from the internal collapses.

Morse theory and Whitehead deformations

Theorem [F.]







Let K be a regular CW-complex of dimension n and let $f : K \rightarrow \mathbb{R}$ be discrete Morse function. Then, f induces a sequence of CW-subcomplexes of K

$$\emptyset = K_{-1} \subseteq L_0 \subseteq K_0 \subseteq L_1 \subseteq K_1 \cdots \subseteq L_N \subseteq K_N = K$$

such that $K_j \searrow L_j$ for all $1 \leq j \leq N$ and $L_j = K_{j-1} \cup \bigcup_{i=1}^{d_j} e_i^j$ with $\{e_i^j : 0 \leq j \leq N, 1 \leq i \leq d_j\}$ the set of critical cells of f . Moreover,

$$K \xrightarrow{\quad n+1 \quad} L_0 \cup \bigcup_{j=1}^N \bigcup_{i=1}^{d_j} \check{e}_i^j = K_{\mathcal{M}}.$$

Example



Morse theory for group presentations

Group presentations \mathcal{P}

Group presentations

\mathcal{P}

CW-complexes

$K_{\mathcal{P}}$



Group presentations

\mathcal{P}

CW-complexes

$K'_{\mathcal{P}}$



Group presentations

\mathcal{P}



CW-complexes

K'_P
+
 f

Group presentations

\mathcal{P}



CW-complexes


K'_P
+
 f



$(K'_P)_M$

Group presentations


CW-complexes



Pipeline

Group presentations


CW-complexes




Pipeline

Group presentations

CW-complexes






We have an algorithmic description of $\mathcal{P}_{(K'_P)_M}$ from \mathcal{P} .

Example

$$\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$$


Example

$$\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$$




Example

$$\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$$



K'_P




+ $f : K'_P \rightarrow \mathbb{R}$ discrete Morse function

Example: $\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$

$\mathcal{Q}_0 = \langle x_2, x_3, x_4, x_5, x_6, \dots, x_{15} \mid x_2, x_3, x_4, x_5,$

$x_7 x_2^{-1} x_5^{-1}, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9 x_2^{-1}, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11} x_3, x_{12} x_{11}^{-1} x_5, x_{13} x_3 x_4^{-1}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15} x_3, x_4 x_{12} x_{15}^{-1} \rangle$




Example: $\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$

$$Q_0 = \langle x_2, x_3, x_4, x_5, x_6, \dots, x_{15} \mid x_2, x_3, x_4, x_5,$$

$$x_7 x_2^{-1} x_5^{-1}, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9 x_2^{-1}, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11} x_3, x_{12} x_{11}^{-1} x_5, x_{13} x_3 x_4^{-1}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15} x_3, x_4 x_{12} x_{15}^{-1} \rangle$$

$$Q_1 = \langle x_6 \dots, x_{15} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15}, x_{12} x_{15}^{-1} \rangle$$




Example: $\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$

$$Q_0 = \langle x_2, x_3, x_4, x_5, x_6, \dots, x_{15} \mid x_2, x_3, x_4, x_5,$$

$$x_7 x_2^{-1} x_5^{-1}, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9 x_2^{-1}, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11} x_3, x_{12} x_{11}^{-1} x_5, x_{13} x_3 x_4^{-1}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15} x_3, x_4 x_{12} x_{15}^{-1} \rangle$$

$$Q_1 = \langle x_6, \dots, x_{15} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15}, x_{12} x_{15}^{-1} \rangle$$

$$Q_2 = \langle x_6, \dots, x_{14} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{12} x_{14}^{-1} \rangle$$



Example: $\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$


$$Q_0 = \langle x_2, x_3, x_4, x_5, x_6, \dots, x_{15} \mid x_2, x_3, x_4, x_5,$$

$$x_7 x_2^{-1} x_5^{-1}, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9 x_2^{-1}, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11} x_3, x_{12} x_{11}^{-1} x_5, x_{13} x_3 x_4^{-1}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15} x_3, x_4 x_{12} x_{15}^{-1} \rangle$$

$$Q_1 = \langle x_6, \dots, x_{15} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15}, x_{12} x_{15}^{-1} \rangle$$

$$Q_2 = \langle x_6, \dots, x_{14} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{12} x_{14}^{-1} \rangle$$

$$Q_3 = \langle x_6, \dots, x_{13} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{12} x_{13}^{-1} \rangle$$



Example: $\mathcal{P} = \langle x, y \mid x^2, xy^{-2} \rangle$

$$Q_0 = \langle x_2, x_3, x_4, x_5, x_6, \dots, x_{15} \mid x_2, x_3, x_4, x_5,$$

$$x_7 x_2^{-1} x_5^{-1}, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9 x_2^{-1}, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11} x_3, x_{12} x_{11}^{-1} x_5, x_{13} x_3 x_4^{-1}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15} x_3, x_4 x_{12} x_{15}^{-1} \rangle$$

$$Q_1 = \langle x_6, \dots, x_{15} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{14}^{-1} x_{15}, x_{12} x_{15}^{-1} \rangle$$

$$Q_2 = \langle x_6, \dots, x_{14} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{13}^{-1} x_{14}, x_{12} x_{14}^{-1} \rangle$$

$$Q_3 = \langle x_6, \dots, x_{13} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{13}, x_{12} x_{12} x_{13}^{-1} \rangle$$

$$Q_4 = \langle x_6, \dots, x_{12} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11} x_{12}^{-1}, x_{12}^2 \rangle$$

$$Q_5 = \langle x_6, \dots, x_{11} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^{-1} x_{11}, x_{11}^2 \rangle$$


$$Q_6 = \langle x_6, \dots, x_{10} \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, x_6^{-1} x_9^{-1} x_{10}, x_{10}^2 \rangle$$

$$Q_7 = \langle x_6, \dots, x_9 \mid x_7, x_6^{-1} x_7^{-1} x_8, x_8^{-1} x_9, (x_9 x_6)^2 \rangle$$

$$Q_8 = \langle x_6, x_7, x_8 \mid x_7, x_6^{-1} x_7^{-1} x_8, (x_8 x_6)^2 \rangle$$

$$Q_9 = \langle x_6, x_7 \mid x_7, (x_7 x_6^2)^2 \rangle$$

$$Q_{10} = \langle x_6 \mid x_6^4 \rangle \sim Q^{**} \mathcal{P}$$



Applications to the Andrews–Curtis conjecture

Theorem [F.]

The following balanced presentations of the trivial group satisfies the Andrews–Curtis conjecture:

- $\mathcal{P} = \langle x, y \mid xyx = yxy, x^2 = y^3 \rangle^\dagger$ [Akbulut & Kirby, 1985]
- $\mathcal{P} = \langle x, y \mid x^{-1}y^3x = y^4, x = y^{-1}xyx^{-1} \rangle$ [Miller & Schupp, 1999]
- $\mathcal{P} = \langle x, y \mid x = [x^{-1}, y^{-1}], y = [y^{-1}, x^q] \rangle, \forall q \in \mathbb{N}$ [Gordon, 1984]

[†]First proved by Miasnikov in 2003 using genetic algorithms.

- *PhD Thesis: X. F., Combinatorial methods and algorithms in low-dimensional topology and the Andrews-Curtis conjecture*, University of Buenos Aires, 2017.
- *Preprint: X. F., 3-deformations of 2-complexes and Morse Theory*, arXiv:1912.00115, 2019 (*new version soon*).
- *Code:*
 - X. F., SageMath Module,
<https://github.com/ximenafernandez/Finite-Topological-Spaces>
 - X. F., Kevin Piterman & Ivan Sadofschi Costa, GAP Package,
<https://github.com/isadofschi/posets>

Work in progress: Computation of persistent fundamental group of point clouds.

email: `x.l.fernandez@swansea.ac.uk`

THANKS FOR YOUR ATTENTION!